Chinese Cabbage 1996

CULTIVAR	Source	Date of 50% Harvest	Mkt. Yield (t/ha)	HEAD CHARACTERISTICS					Bolting %	Head Type	Tip Burn	Overall Rating	Comments
				Weight (kg)	Height (cm)	Diam. (cm)	Petiole Color	Leaf Color					
Akala	EZ	Aug. 14	6.5	0.7	27	12	W	G	0	N	5	1.5	Moderate rot by mid-season, severe tip burn
Banko (Bejo 1683	BJ	Sep. 11	39.1	1.3	46	12	LG	DG	0	M	2	4.5	Excellent yields
Blues	J, Os	Aug. 14	25.8	1.9	29	14	W	G	0	N	5	2.5	Moderate-severe rot by mid-season, prickly leaves
Chi Hi Li	EFG	---	0.0	---	---	---	---	---	40	M	2	1	Bolted prior to harvest
China Express	SAK	Aug. 14	28.0	1.5	28	13	W	DG	0	N	5	3	Severe tip burn
China Pride	ST	Aug. 14	21.4	1.7	29	22	LG	DG	0	N	2	2.5	Moderate-severe rot by mid-season
Early Jade Pagoda	Os	Aug. 14	24.8	1.6	35	12	LG	DG	0	M	2	3.5	Nice shape, severe tip burn late in season
Gracious	KY	---	0.0	---	---	---	---	---	100	PC	---	1	Bolted prior to harvest
Green Petiole	KY	---	0.0	---	---	---	---	---	90	PC	1	1	Bolted prior to harvest
Jade Pagoda	ST	Aug. 14	20.1	1.3	37	13	LG	G	5	M	2	3.5	Nice cultivar, attractive
Joi Choi	Ve, CA, ST	Aug. 14	24.5	1.1	37	10	W	DG	0	PC	2	3	Nice cultivar, moderate bolting and severe rot late in season
Kasumi	ST	Aug. 14	40.9	1.4	27	17	W	G	0	N	3	3.5	High yields, attractive but internal insect damage
Lei Choi	AS	Aug. 14	13.2	0.6	36	8	LG	DG	10	PC	3	2	Severe tip burn and 100% bolting late in season, loose heads
Manoko (Bejo 168	BJ	Aug. 14	27.1	1.5	28	14	LG	G	0	N	3	3.5	Good yields, resistant to bolting
Mei Qing Choi	Os	Aug. 14	7.4	0.6	25	8	LG	DG	0	PC	1	1.5	100% bolting late in season
Michihili	FG, TERR, BU	---	0.0	---	---	---	---	---	55	M	1	1	Poor germination, bolted prior to harvest
Ming Choi	ST	Aug. 27	25.7	0.9	40	9	W	DG	15	PC	1	2	Severe rot by mid-season, loose heads, multi-headed plants
Monument	ST	Aug. 14	24.0	1.6	40	26	LG	DG	5	M	2	4	Firm heads, nice cultivar
Nerva	EZ	Aug. 8	36.1	2.2	29	14	W	LG	0	N	2	3.5	Early, nice cultivar, 100% bottom rot by late season, big leaf veins
No. 534 (Crispy)	EG	---	0.0	---	---	---	---	---	100	N	---	1	Bolted prior to harvest
No. 539 (Beau)	EG	---	0.0	---	---	---	---	---	100	N	---	1	Bolted prior to harvest
No. 541 (Gloria)	EG	---	0.0	---	---	---	---	---	100	N	---	1	Bolted prior to harvest
One Kilo SB	TOK	Aug. 14	28.6	1.4	26	10	W	G	0	N	2	3.5	Good yields, resistant to bolting
Optiko	BJ	Aug. 14	43.4	1.7	28	17	W	G	0	N	2	3.5	High yields, moderate-severe rot late in season
Pak Choi	EFG	Aug. 14	56.3	0.9	44	9	LG	G	0	PC	2	4.5	Nice cultivar but loose heads, minimal insect damage, high yields
Shori	NH	Aug. 14	36.5	1.5	25	18	W	G	0	N	3	3	Good yields, moderate-severe rot late in season, insect damage
Spring Flavor	HN	Aug. 14	16.1	1.1	24	15	LG	DG	0	N	3	2.5	Moderate-severe rot late in season
Summer Top	J	Aug. 14	4.1	0.7	23	7	LG	DG	0	N	4	1.5	Pick when small, bolts by late season
TSX-4111	TOK	Aug. 14	10.7	0.8	21	9	LG	G	0	N	5	2	Severe tip burn
TSX-4112	TOK	Aug. 27	7.4	1.1	22	9	W	G	0	N	3	2	Severe rot late in season
White Sun	KY	---	0.0	---	---	---	---	---	100	N	---	1	Bolted prior to harvest
WR70	Os	Aug. 14	27.2	1.3	26	15	W	G	0	N	2	2.5	Severe tip burn late in season, insect damage
Yuki	SAK	Aug. 14	30.4	1.3	25	14	W	G	0	N	2	2.5	Severe tip burn late in season, insect damage

Date seeded: May 28

Spacing: In row - 15 cm
Between row - 75 cm

Head type: M - Michihili
N - Napa
PC - Pak Choi

Color: G - green
LG - light green
DG - dark green
W - white

Tip burn: 1 - minimal
5 - severe

Overall rating: 1 - poor
5 - excellent

Bolting % and Tip Burn ratings were recorded July 30.